

Nuclear and Particle Physics applications of the Bohm Picture of Quantum Mechanics

A. Miranda

January 9, 2009

Department of Physics and Astronomy, Aarhus University, DK-8000
Aarhus C, Denmark

Abstract

Aproximation methods for calculating individual particle/ field motions in spacetime at the quantum level of accuracy (a key feature of the Bohm Picture of Quantum Mechanics (BP)), are studied.

This sharply illuminates not only the deep quantum structures underlying any observable quantum statistical laws of motion of particles and fields in spacetime, but also how the continuous merging of the so-called classical and quantal modes of description actually occurs, with no breaks anywhere.

Modern textbook presentations of Quantum Theory are used throughout, but only to provide the necessary, already existing, tested for-

malisms and calculational techniques. New coherent insights, reinterpretations of old solutions and results, and new (in principle testable) quantitative and qualitative predictions, can be obtained on the basis of the BP that complete the standard type of postdictions and predictions.

Most of the dead wood still cluttering discussions on the meaning of Quantum Theory and the role of the BP is by-passed.

We shall try to draw attention to the physics of this unfortunately hardly known novel formulation of Quantum Theory by giving additional illustrative examples inspired from the daily practices of contemporary Nuclear and Particle Physics, subjects that as yet have not been thoroughly reinterpreted within the BP.

These fields of research offer excellent opportunities for explaining and illustrating the significance of time in quantum transitions, as well as the closely related features of quantum non-locality and quantum wholeness, as hard physical facts. We claim that in addition we can obtain a substantial gain in predictive powers of the underlying, all-encompassing, Quantum Theory.

PACS numbers: 03.65, 23.40.-s, 23.60+e, 28.20-v

1 A short introduction to the Bohm Picture of Quantum Mechanics (BP)

Quantum Theory, as it is explained in textbooks, comes in two fundamental versions, classified by Dirac with the word *Pictures* [1]: the Schroedinger Picture (SP), where time developments are entirely accounted for by the wavefunctions (which can be described in many different *representations*) and the Heisenberg Picture (HP), where time-developments are entirely encoded in the operators.

1.1 Starting from the SP

The differential form of the Schroedinger equation of motion is (we set $\hbar = 1$, unless otherwise stated):

$$i \frac{d}{dt} |\Psi(t)\rangle = H |\Psi(t)\rangle \quad (1)$$

An integral representation of the Schroedinger ket vector $|\Psi(t)\rangle$ is

$$|\Psi(t)\rangle = e^{-iH(t-t')} |\Psi(t')\rangle \quad (2)$$

The Hamiltonian H in this context has a double role: it is the operator generating time-translations, assumed to be itself time-independent (for isolated systems), and it is the operator whose eigenvectors are stationary states Φ_ν of the system:

$$H(N)\Phi_\nu(\{\vec{x}\}; E_\nu) = E_\nu\Phi_\nu(\{\vec{x}\}; E_\nu) \quad (3)$$

We shall use the simplifying short-hand notation:

$$\{\vec{x}\} \equiv \vec{x}_1, \vec{x}_2, \dots, \vec{x}_N \quad (4)$$

$$\{\vec{p}\} \equiv -i\vec{\nabla}_1, -i\vec{\nabla}_2, \dots, -i\vec{\nabla}_N \quad (5)$$

In general, let us assume some complete set of state vectors in Hilbert space:

$$\sum_{\alpha} |\alpha\rangle\langle\alpha| = 1 \quad (6)$$

$$\langle\beta|\alpha\rangle = \delta(\beta, \alpha) \quad (7)$$

Then the wavefunctions (or their complex conjugates) are just components (complex numbers) of the ket vector $|\Psi(t)\rangle$ (or the bra vector $\langle\Psi(t)|$) in the Hilbert space in whatever the chosen basis vectors is [1] :

$$\Psi_\beta(t) = \langle\beta|\Psi(t)\rangle \quad (8)$$

$$\Psi_\alpha^*(t) = \langle\Psi(t)|\alpha\rangle \quad (9)$$

We work in this paper with an entirely different Picture, unfortunately not even mentioned in textbooks. To give it a name, we shall refer to it as the Bohm Picture (BP)[2,3,4] and shall be discussed presently.

Before we proceed, let us ask the question : would it make any real difference, had Bohm started instead from the HP, instead of the SP?

From one of Dirac's last publications [1] we can infer that for non-relativistic systems the answer is NO; for relativistic Quantum Field Theory (QFT), the case seems to remain open.

Practical approximation methods for solving the time-dependent and time-independent Schroedinger equation, for realistic systems, both in the textbook formulation and in published BP papers [2,3], are generally based on some standard combinations of Perturbation - Variation theory with computer power.

As we shall see, wavefunctions associated with specific physical systems are an essential part of input information, but only a part. This statement sharply disagrees not only with textbook quantum theory but also with all (as far as I know) alternative ontological formulations [2].

Let us attempt to look more closely into the question of time development in Quantum Mechanics, as accounted for in the BP. As it is perhaps widely agreed, an universal time seems to be required by the very definition of the Schroedinger equation.

We follow Bohm by choosing the basis set of wavefunctions to be the

eigenstates of the Schroedinger (hermitean) “position operators” $\hat{x}_k$:

$$\hat{x}_k|\vec{x}_1, \vec{x}_2, \dots, \vec{x}_N\rangle = \vec{x}_k|\vec{x}_1, \vec{x}_2, \dots, \vec{x}_N\rangle \quad (10)$$

$$[\hat{x}_k, \hat{x}_j] = 0 \quad (11)$$

$$[\hat{p}_k, \hat{p}_j] = 0 \quad (12)$$

$$[\hat{x}_k, \hat{p}_j] = i\delta(k, j) \quad (13)$$

$$\Psi(\{\vec{x}\}; t) \equiv \langle \{\vec{x}\} | \Psi(t) \rangle \quad (14)$$

The time-dependent Schroedinger equation for describing a system with countable $N=1,2,\dots$, degrees of freedom can always be rewritten in the form

$$H(N)\Psi(\{\vec{x}\}; t) = i\frac{\partial}{\partial t}\Psi(\{\vec{x}\}; t) \quad (15)$$

where $H(N)$ is the (supposed hermitean) Hamiltonian:

$$H(N) = U_0 - \sum_{i=1}^N \frac{1}{2m_i} \nabla_i^2 + V(\{\vec{x}\}) \quad (16)$$

U_0 is an arbitrary constant. The function V represents a *preassigned*

classically definable *local* function:

$$< \{\vec{x}'\} | V | \{\vec{x}\} > \equiv \delta(\{\vec{x}'\} - \{\vec{x}\}) V(\{\vec{x}\}) \quad (17)$$

So a general time-dependent wavefunction is obviously

$$\Psi(\{\vec{x}\}; t) = \sum_{\nu, E_\nu} C_\nu(E_\nu) \Phi(\{\vec{x}\}; E_\nu) \exp(-iE_\nu t) \quad (18)$$

and is completely defined, if the values of the coefficients C_ν be given at some initial time $t=0$:

$$\Psi(\{\vec{x}\}; 0) = \sum_{\nu, E_\nu} C_\nu(E_\nu) \Phi(\{\vec{x}\}; E_\nu) \quad (19)$$

Thus assuming that an initial wave-packet is given, we immediately deduce an explicit expression for the coefficients

$$\Psi(\{\vec{x}\}; 0) = \sum_i a_i \varphi_i(\{\vec{x}\}) \quad (20)$$

$$\Phi_\nu(\{\vec{x}\}, E_\nu) = \sum_i b_{\nu i}(E_\nu) \varphi_i(\{\vec{x}\}) \quad (21)$$

$$C_\nu(E_\nu) = \sum_i b_{\nu i}^*(E_\nu) a_i \quad (22)$$

where Ψ and Φ are expanded over some convenient complete set of functions $\{\varphi_i\}$.

We shall simply assume familiarity with this standard material and carry on from there. Let us next introduce the BP on the basis of the following

postulates:

POSTULATE 1

As the wavefunction is in general a complex number, let us write

$$\Psi(\{\vec{x}\}; t) = R(\{\vec{x}\}; t) e^{iS(\{\vec{x}\}; t)} \quad (23)$$

where R and S are both real.

This by definition a *beable* [2,3,4] officially known “wavefunction” . It obeys its own equation of motion, which is none other than the Schroedinger equation. The latter can be rewritten as a system of time-dependent coupled differential equations involving two real functions R and S:

$$\begin{aligned} & \frac{\partial S(\{\vec{x}\}; t)}{\partial t} + \\ & + \sum_{k=1}^N \frac{1}{2m_k} |\vec{\nabla}_k S(\{\vec{x}\}; t)|^2 + V(\{\vec{x}\}) + Q(\{\vec{x}\}; t) \end{aligned} \quad (24)$$

$$\frac{\partial P(\{\vec{x}\}; t)}{\partial t} + \vec{\nabla} \cdot \sum_k \vec{J}_k(\{\vec{x}\}; t) = 0 \quad (25)$$

with the definitions

$$\vec{J}_k(\{\vec{x}\}; t) = \vec{\nabla}_k S(\{\vec{x}\}; t) \quad (26)$$

$$P \times \vec{J}_k = Im[\Psi^* \vec{\nabla}_k \Psi] \quad (27)$$

$$P = R^2 \tag{28}$$

$$Q(\{\vec{x}\}; t) \equiv - \sum_k \frac{1}{2m_k} \frac{\nabla_k^2 R(\{\vec{x}\}; t)}{R(\{\vec{x}\}; t)} \tag{29}$$

The last quantity introduced is known as the " Quantum Potential" or rather, "Quantum Information Potential" (QIP) [2] and it is, together with definition (26), the core of the alternative formulation of non-relativistic quantum mechanics proposed by Bohm and his co-workers [2,3,4]. As shown by these authors, it can be naturally generalised to Quantum Field Theory (QFT).

In any case, we emphasize that all the above definitions are derived from the time-dependent Schroedinger equation, and that their fundamental status stands or falls with the validity of this equation.

Let us first introduce the other set of further relevant *beables* [2,4] in BP, i.e. particles and fields: this *completes* the formulation of Quantum Theory in the BP.

One begins by defining dynamical quantities such as particle momenta, in complete agreement - in the appropriate limit of course, i.e. when Q is negligible - with the classical definition of momenta that is part of the Hamilton-Jacobi formulation of classical mechanics [2,3]. We thus get Bohm's equations of motions for particles

$$\begin{aligned}
\vec{p}_k(\{\vec{X}(t)\}; t) &= m_k \frac{d\vec{X}_k(\{\vec{X}(t)\}; t)}{dt} = \\
&= \{\vec{\nabla}_k S(\{\vec{x}\}; t)\}_{\{\vec{x}\}=\{\vec{X}(t)\}}
\end{aligned} \tag{30}$$

or (equivalently) the velocity of a particle k at its position $\vec{X}_k(t)$:

$$\begin{aligned}
&\frac{d\vec{X}_k(\{\vec{X}(t)\}; t)}{dt} = \\
&= \frac{1}{m_k} \left[\frac{1}{P} \text{Im}(\Psi^* \vec{\nabla}_k \Psi) \right]_{\{\vec{x}\}=\{\vec{X}(t)\}}
\end{aligned} \tag{31}$$

This non-linear set of ordinary differential equations is the fundamental set of Bohm equation of motion for *beable* particles [2]: it is the essential part of the theoretical framework that, according to Bohm and co-workers, is missing from standard textbook formulations and alternative ontological interpretations [2]. It is then demonstrated that a complete removal of all the dead wood of ambiguities and “mysteries” of textbook QM can thereby be elegantly achieved.

Eq.(28) can be rearranged to give an alternative version:

$$m_k \frac{d^2 \vec{X}_k(t)}{dt^2} =$$

$$= -[\vec{\nabla}_k(V(\{\vec{x}\}) + Q(\{\vec{x}\}; t))]|_{\{\vec{x}\}=\{\vec{X}(t)\}} \quad (32)$$

One can thus clearly see that any beable particle is subject to *quantum accelerations*, regardless of the existence of classical (inertial) accelerations.

This sharply illuminates the profound difference between what a “particle” means in classical physics and what a “beable particle” means in Bohm’s quantum physics [2]. Unawareness of, or misunderstandings about this fundamental difference can only lead to unnecessary confusions and endless irrelevant objections and production of paradoxes.

The individual Bohm beable particle history (i.e. existing regardless of whether the particle is, or is not, a member of an ensemble) is then formally defined to be a solution of these non-linear ordinary differential equations (31),(32).

Key features of quantum theory, such as quantum non-locality and quantum wholeness [2], are thus precisely expressed by these equations for particle (and field) motions as they are ultimately interpreted in spacetime:

(i) single particle histories are primarily completely entangled (i.e. *in principle* defined by, unseparable from) the entire environment (literally, the entire universe), because they are *actively quantum in – formed* by the *phase* of the wavefunction [2]. This has of course nothing to do with *classically* defined “medium entanglement” (many-body effects) through “potentials, gravitation, thermal influences” and so forth;

(ii) an essential feature of this quantum information is the de Broglie *guidance* condition on the particles eq (30), which puts the *potential* beable information encoded in the common wave function into the *actual* beable motion of every *individual* beable particle (as distinct from statistical ensembles of such particles), participating in the potentially present common pool of information provided by the wavefunction [2].

So, the physics at the quantum level of accuracy differs totally from that assumed and expressed by classical mechanics, first and foremost because it is this *quantum wholeness* that appears to be the primary reality [2]. This is encoded *abinitio* in the origin, structure and every feature of the QIP [2].

It should be quite clear that one is indeed introducing here crucial novel concepts that altogether single out the BP.

The beable Bohm histories of particles (and fields) then go *continuously* over to *observable* newtonian (or Hamilton-Jacobi) trajectories. But one should be in no doubt [2] that the presence of the QIP profoundly changes the entire situation from the bottom up, in a way that can be neither understood nor interpreted in classical mechanics terms. Thus the emergence of familiar classical spacetime descriptions is indeed an immediate consequence of certain especial features of the QIP.

Two extreme situations can be immediately envisaged:

(i) The QIP is numerically completely negligible relative to the classical potential (or, in more physical terms, compared to unavoidable thermal fluctuations). In this case, the quantum forces derived from it are completely

overrun by ordinary classical mechanical and thermodynamical forces.

A *continuous transition* to a classical regime (i.e. pure spacetime descriptions, with use of coordinates) then occurs. Numerical quantum corrections are then too small to play any important role.

(ii) The QIP is separable, i.e.

$$Q(\{\vec{x}\}; t) =$$

$$= - \sum_{k=1}^N \frac{1}{2m_k} \frac{1}{R_k(\{\vec{x}\}; t)} \nabla_k^2 R(\{\vec{x}\}; t) \equiv \sum_{k=1}^N Q_k(\{\vec{x}\}; t) \quad (33)$$

In this case, the specific quantum mechanical non-locality dissolves and each particle behaves classically as if completely unaware of its surroundings, except that it may interact with these surroundings through eventual classically definable preassigned potentials, or more physically speaking, through the omnipresent thermal fluctuations.

POSTULATE 2:

In order to confront the BP (one should never forget that BP deals primarily with individual beables [2,3,4]) with any existing experimental material, one needs first to analyse in detail the anatomy of actual measurement processes.

This was thoroughly done by de Broglie, Bohm and their co-workers [2,3,4] in general, at least for non-relativistic systems. The situation when relativistic requirements are essential is still far less digested, both in text-

book QFT and in the relativistic BP [2,3,4].

Let us then simply summarize one of the main results : one has to apply some definite stochastic averaging procedures to the above theory of individual beables, very much the same way as it is and was always done in a classical context, e.g. in classical Statistical Mechanics [2,3,4].

Given these individual Bohm histories of beables, it is straightforward to compute statistical averages (i.e. averaging all initial positions and velocities) over pure ensembles of such histories [2,3] . For example, assuming for simplicity just one spatial dimension,

$$\langle X(t) \rangle_{EA} \equiv \int dX(t) R^2(X(t), t) X(t) \quad (34)$$

$$\langle P(t) \rangle_{EA} \equiv \int dX(t) R^2(X(t), t) \left[\frac{\partial S(x, t)}{\partial x} \right]_{x=X(t)} \quad (35)$$

$$\langle E(t) \rangle_{EA} \equiv \int dX(t) R^2(X(t), t) \left[-\frac{\partial S(x, t)}{\partial t} \right]_{x=X(t)} \quad (36)$$

where the history $X(t)$ is given by (30). This turns out to be - of course! - exactly the same as ordinary quantum mechanical expectation values, taken with the *same* wavefunction.

So, statistics (a practical necessity, simply because *we* cannot know, nor is it necessary *for us* to know, everything that happened, happens and shall happen in the world!) does not have in the context of the present formulation the same fundamental status given to it by textbook formulations of quantum

theory, which deal solely with “observables”, i.e. with the actual experimental data resulting from actually performed experiments.

Most of these observables are context dependent, because of the participatory nature of “observations” in Quantum Mechanics [2]. Perhaps the one important exception to this fact of life is the observable “position of things” [4], i.e the eigenvalues of the spatial position operator. This observable can be directly linked to the beable “position of (discrete) particles in space” or “continuous fields at specified points in space”, introduced in the previous postulate. Space is, after all, where we exist and do experiments! It is the playground of classical physics (and of common sense).

As both de Broglie and Bohm demonstrated, both mathematically and using physics, the probability distributions to do this necessary stochastic averaging, whatever they might originally be, rapidly relax to none other than the familiar Born probabilities [2]. Because of the local conservation of probability (25), if this holds at some particular time, it will hold for all times. Hence, within the context of any given real experiment, the results predicted in principle by textbook and the ensemble averaged BP over all initial positions *and* velocities must be exactly the same.

A misunderstanding (or unawareness) of this most important POSTULATE 2 has caused a great deal of unnecessary confusion in the published literature [5] about the meaning and status of the BP.

Real differences would be seen only for histories of individual systems, as distinct from ensembles of such systems. Only if, and when, someone some

day has gathered sufficient such data can one make meaningful judgements on the status of BP.

2 Case studies from contemporary Nuclear and Particle physics

We consider simple, but illustrative, additional examples inspired by contemporary practices in Nuclear and Particle Physics.

These fields of research offer excellent new opportunities for testing specific interpretations and predictions based on the BP of Quantum Theory. A good reason is, they often draw attention to the understanding of two key features of all (quantum) processes, viz. the question of transitions in time and the closely related feature of quantum non-locality.

2.1 EPR-like correlations in pure α -decay channels of very light nuclei

Among the very light nuclei, the ^8Be and ^{12}C isotopes can decay into pure two or three α -channels with appreciable branching ratios. These channels always played an important role in the theories and experiments on ordinary stellar evolution. Thus the momenta and energy distributions of the final state particles have been thoroughly investigated [6,7,8], but not the predictable EPR-like quantum correlations - the main issue here. The point of

interest is that the outgoing α 's in these decays are bosons, even if the parent nuclei are bags of nucleons, which are fermions. So the wavefunctions in the outgoing decay channels must be fully symmetric under exchanges of arbitrary labels attached to them (in this case, just their position coordinates). So, any model candidate to a beable wavefunction in the Center of Mass (CM) reference frame of this nucleus must be a fully symmetric function of the internal coordinates.

We shall ignore for simplicity the final state Coulomb interaction [7] that in principle operates among these outgoing particles and accelerates them away from each other. We could assume, as it is done in actual numerical fits [7], that this Coulomb interaction is screened; in any case, it plays only a secondary role here, as it leads basically to what could be called *classical entanglement*.

However, the outgoing α -channels, even if not interacting through pre-assigned classical potentials, are EPR-correlated through the Bose symmetry requirement on the wavefunction [2]. As is well-known these EPR-correlations are essentially non-local [2], and thus beyond any classical accountability.

We shall show from the viewpoint of the BP of Quantum Mechanics how these correlations (that must be there) can be traced back to the symmetry requirement on the wavefunction and, as a by-product, attempt to give a precise meaning to words such as “identical, equivalent, distinguishable, undistinguishable,...” particles.

In order to make this discussion as simple as possible but without losing generality, let us consider only one spatial dimension and imagine a ${}^8\text{Be}$ nucleus (at rest when $t=0$) fissioning into two α -particles. Extensions to heavier nuclei decays involving more α - channels [6] do not add essentially new physics, but do complicate the detailed mathematics considerably.

So, let the Hamiltonian for $t>0$ be

$$H = -\frac{1}{2M_\alpha} \sum_{k=1}^2 \frac{\partial^2}{\partial x_k^2} \quad t > 0 \quad (37)$$

A central feature of this Hamiltonian is its invariance under the two permutations of the labels 1,2 as a result of a common mass for the particles. True stationary states must then share this symmetry with the Hamiltonian. Quantum Mechanics establishes that these must be fully symmetric functions of the coordinates, as one is dealing with bosons [10]. The Schroedinger equation is

$$H\Psi(x_1, x_2; t) = i\frac{\partial}{\partial t}\Psi(x_1, x_2; t) \quad t > 0 \quad (38)$$

It must be complemented with the initial condition

$$\Psi(x_1, x_2; 0+) = F(x_1, x_2) \quad (39)$$

where F is a given (fully symmetric) function of spatial coordinates. It is assumed of course that the wavefunction is continuous at $t=0$.

The consequences of the symmetry requirement are obvious, if we trans-

form to Jacobi coordinates:

$$x = x_1 - x_2 \quad (40)$$

$$X_{CM} = \frac{x_1 + x_2}{2} \quad (41)$$

The coordinate x is antisymmetric under the exchange $1 \leftrightarrow 2$ whereas X_{CM} is symmetric. Hence we must have,

$$\Psi(x_1, x_2; t) = g(x; t) \times e^{iPX_{CM}} \quad (42)$$

where

$$g(x; t) = g(-x; t) \quad (43)$$

Let us assume that we are in the CM coordinate system First consider a single particle guided by the wave-packet

$$\begin{aligned} \Psi(x, t; \{A\}) &= \int_{-\infty}^{+\infty} \exp(-i \frac{p^2}{2M_\alpha} t) * e^{-(p-p_A)^2/\sigma_A^2} * \\ &* \exp(ip(x - x_A)) dp \quad t > 0 \end{aligned} \quad (44)$$

Putting

$$\Psi(x, t; \{A\}) = R(x, t; \{A\}) \exp(iS(x, t; \{A\})) \quad (45)$$

we find that the particle velocity at time t is given by the guidance condition (30):

$$\begin{aligned} \frac{dX(t; \{A\})}{dt} &= \frac{1}{M_\alpha} \left\{ \frac{\partial}{\partial x} S(x, t; \{A\}) \right\}_{x=X(t)} = \\ &= \frac{1}{1 + (\frac{\sigma_A^2}{2M_\alpha} t)^2} \left[\frac{p_A}{M_\alpha} + \frac{\sigma_A^2}{M_\alpha} (X(t; \{A\}) - x_A) t \right] \end{aligned} \quad (46)$$

A simple quadrature then gives the particle position at time t . This shows that the particle is accelerated from its initial velocity p_A/M_α at time $t=0$ to its final velocity at $t=+\infty$.

The fundamental reason for this pure quantum effect is, as discussed in Section 1, the action of the QIP(29) that is directly linked to the Schroedinger wavefunction.

Next, consider two α 's but ignore the symmetry requirement. Then the wavefunction is simply

$$\Psi(x_1, x_2, t; \{A, B\}) = \Psi(x_1, t; \{A\}) \times \Psi(x_2, t; \{B\}) \quad (47)$$

hence

$$R(x_1, x_2, t; \{A, B\}) = R(x_1, t; \{A\}) \times R(x_2, t; \{B\}) \quad (48)$$

$$S(x_1, x_2, t; \{A, B\}) = S(x_1, t; \{A\}) + S(x_2, t; \{B\}) \quad (49)$$

Using the guidance condition (30) the velocities are

$$\frac{d}{dt}X_1(t; A) = \left\{ \frac{\partial}{\partial x_1} S(x_1, t; \{A\}) \right\}_{x_1=X_1(t)} \quad (50)$$

$$\frac{d}{dt}X_2(t; B) = \left\{ \frac{\partial}{\partial x_2} S(x_2, t; \{B\}) \right\}_{x_2=X_2(t)} \quad (51)$$

We conclude that the α -particle that was labelled 1 at time $t=0$ follows the beable trajectory A, i.e. $X_1(t; A)$, and is therefore still the same particle at any later time t , exactly as one classically would expect ; the same applies to particle 2, moving along the beable spacetime trajectory $X_2(t; B)$. As expected classically, 1 and 2 are completely “unaware” of each other, in spite of the fact that they have a common origin, that is, the fissioning of ${}^8\text{Be}$. There is no question about which is 1 and which is 2, as one can always in principle follow their trajectories at any time. This applies, even if 1 and 2 are “indistinguishable, equivalent, ...”. We are so far in complete harmony with classical physics thinking.

Next, consider again two α 's but insist that the symmetry requirement must be obeyed by the wavefunction, as it should. Then the correct wavefunction is

$$\begin{aligned}
\Psi(x_1, x_2, t; \{A, B\}) &= \frac{1}{\sqrt{2}} \{ \Psi(x_1, t; \{A\}) \times \Psi(x_2, t; \{B\}) + \\
&\quad + \Psi(x_2, t; \{A\}) \times \Psi(x_1, t; \{B\}) \} \\
&= \Psi(x_2, x_1, t; \{A, B\})
\end{aligned} \tag{52}$$

Now the separability expressed by the definitions (48) and (49) is lost, and the results (50) and (51) are no longer valid. Then the more general condition (30) prevails:

$$\begin{aligned}
\frac{d}{dt} X_1(t; A, B) &= \\
&= \left\{ \frac{\partial}{\partial x_1} S(x_1, x_2, t; \{A, B\}) \right\}_{x_1, x_2 = X_1(t), X_2(t)}
\end{aligned} \tag{53}$$

$$\begin{aligned}
\frac{d}{dt} X_2(t; A, B) &= \\
&= \left\{ \frac{\partial}{\partial x_2} S(x_1, x_2, t; \{A, B\}) \right\}_{x_1, x_2 = X_1(t), X_2(t)}
\end{aligned} \tag{54}$$

This is tantamount to quantum non-locality, here meaning dependence on $x = x_1 - x_2$ that persists no matter how large x is, even in absence of any classically describable forces.

So, a natural (and classically meaningful) question would be: suppose one places an α -detector at a position D_A and simultaneously another identical α -detector at a position D_B . These detectors duly detect at some time t the two α -particles upon their arrival there. But is the particle that is detected at D_A the “same” particle called 1 at time $t=0$, and the particle detected at D_B the “same” called 2 at time $t=0$? The exact and real answer is given by (53) and (54) which, within the classical physics philosophy (and within the philosophy of the usual common language as well), are meaningless!

Another feature of results (53) and (54) that are closer to the original EPR-correlation problem, is: instead of position detectors we could try momentum detectors. Eqs.(53)and (54) show that the *momentum* of say particle 1 at time t must depend on the *position* at the *same time t* of particle 2, independently of the distance separating the two particles and vice-versa. Note that this conclusion applies to individual beables and thus has nothing to do with the usual probabilities and statistics. However, upon taking appropriate ensemble averages (Section 1), i.e. doing the usual statistics of measurements [2,3,4], we would end up exactly with the standard textbook predictions, which apply of course only to some actual experimental result and refer only to that particular experimental set up that produces that result.

2.2 Neutron interferometry under gravitational fields

The subject of specific quantum features seen in accelerated reference systems is a promising field for future experimental research. Generally speaking, ultracold neutron physics [9] is especially interesting, due to high-quality experimental work now available.

Essentially quantum effects are deeply linked with changes in the phase of the wavefunction [10]. So, the most revealing tell-tale of the underlying quantum reality is most simply and directly provided by interference experiments. Classical examples are provided by the Aarhanov-Bohm and the Aarhanov-Casher effects [2,3].

More recently, very precise experimental results on motions of neutrons in the earth's gravitational field became available [9]. Consider a free neutron in motion under the gravitational field of the earth [9,10]. Classically, any massive particle would move along the newtonian trajectory given by a solution of the equation of motion

$$m_{in} \frac{d^2 z}{dt^2} = -m_{gr} g \quad (55)$$

where the z-axis is chosen to be parallel to the gravitational lines of force. As well-known [11]

$$m_{in}/m_{gr} = \textit{universal constant} \quad (56)$$

So a choice of mass units allows this universal constant to be set to unity,

and we get a simple *geometrical* law of motion under gravity

$$\frac{d^2 z(t)}{dt^2} = -g \quad (57)$$

$$z(t) = z(0) - \frac{1}{2}gt^2 \quad (58)$$

quite independent of the mass parameter.

When using QM to reexamine this phenomenon, this most remarkable independence on mass seems to disappear. However, this holds only if $\hbar \neq 0$, which of course is the case: the Schroedinger equation of motion is then

$$\left[-\frac{d^2}{dz^2} - 2\left(\frac{M_N}{\hbar}\right)^2 gz\right]\Psi(z, t) = 2i\left(\frac{M_N}{\hbar}\right)\frac{\partial}{\partial t}\Psi(z, t) \quad (59)$$

where M_N is the neutron mass. So the mass does contribute, but only in the combination $M_N/\hbar$.

Note that there is no contradiction at all with the experimental fact [11], just as there isn't any in classical mechanics.

Thus the Bohm equation of motion for any individual neutron that replaces (57) can be written down. The neutron velocity is given by eq

$$v(Z(t), t) = \frac{1}{M_N} \left[\frac{d}{dz} S(z, t) \right]_{z=Z(t)} \quad (60)$$

where $S(z, t)/\hbar$ is the phase of the wavefunction $\Psi(z, t)$. From this the neutron history can be predicted, given the initial conditions.

Consider now the following experiment [10]: a neutron beam with energy E is split into two channels, a channel ABD and another channel ACD. The channels are made to meet again at the neutron detector D. For example, in the table-top experiment discussed in [10] the neutron following the path ABD first climbs from the ground level up an inclined plane (say, at angle δ with the ground) from A to B along a path of length l_1 then at B follows the path BD (length l_2) parallel to the ground (thus normal to the gravitational lines of force) until it reaches the detector at D. If a neutron is instead guided along the path ACD, the opposite sequence is followed, i.e. first AC (corresponding to BD) then CD (corresponding to AB) until the neutron reaches the detector at D. The detectors at D record the arrival at D of any neutron member of a statistical ensemble, sharing the same wavefunction, as a function of the inclination δ for a given beam energy E .

This is of course just another example of a “which-path” experiment, this time using neutron interferometry [10]. Let us then assume that

$$\begin{aligned}\Psi(Z(t), t) = & R_1(Z(t), t) \exp(iS_1(Z(t), t)/\hbar) + \\ & + R_2(Z(t), t) \exp(iS_2(Z(t), t)/\hbar)\end{aligned}\tag{61}$$

The probability pattern at the detector is then

$$P(Z(t), t) = R_1^2(Z(t), t) + R_2^2(Z(t), t) + 2R_1R_2\cos\Phi_{12}\tag{62}$$

where Φ_{12} is the phase difference between the two paths :

$$\Phi_{12} = (S_1(ABD) - S_2(ACD))/\hbar \quad (63)$$

It can be quickly estimated in the WKB approximation (which, remember, neglects the QIP). Phase change along ABD:

$$\frac{S_{ABD}}{\hbar} = \int_{ABD} p dl / \hbar = \int_{ABD} \sqrt{2M_N(E - M_N g z)} dl / \hbar \quad (64)$$

and along ACD:

$$\frac{S_{ACD}}{\hbar} = \int_{ACD} p dl / \hbar = \int_{ACD} \sqrt{2M_N(E - M_N g z)} dl / \hbar \quad (65)$$

hence

$$\Phi_{12} = -l_2 \sqrt{2M_N E} / (2E\hbar) * M_N g l_1 \sin \delta \quad (66)$$

Recalling that

$$\sqrt{2M_N E} / E = 2M_N \lambda / \hbar \quad (67)$$

where λ is the de Broglie wavelength of the incoming wavefunction. We thus get our final result [10]

$$\Phi_{12} = -(M_N / \hbar)^2 g l_1 l_2 \sin \delta * \frac{\lambda}{2\pi} \quad (68)$$

This prediction appears to be in excellent agreement with the experimental results [9,10].

Let us now consider the realistic case that any incoming neutron with given velocity belongs in a statistical ensemble of such particles.

Upon arrival at time $t=0$ at A they are all redirected (i.e. the trajectories bifurcate). Depending on the initial position and velocity of a particle, either the path ABD or the path ACD will be followed, but every single body is forced to meet at D.

The histories $Z(t)$ of individual neutrons are given by formula (30). The actually observed pattern of arrivals of individuals at D is fixed by (62).

The *probability* that at time $t=0$ any particular neutron *is* at $Z=0$ with velocity $V(0)$ is *assumed* (Postulate 2) to be given by

$$P(Z(0) = 0, t = 0) = |\Psi(Z(0) = 0, t = 0)|^2 \quad (69)$$

We emphasize that in BP there is nothing fundamental about this assumption.

As a matter of fact, this is the only place where probabilities enter the game at all, just as they do in Classical Statistical Mechanics. Everything else for $t>0$ is completely determined by the initial conditions. Thus the probability that a neutron *will be* at the position $Z(t)$ (with velocity $V(t)$ given by (30)) is

$$P(Z(t), t) = |\Psi(Z(t), t)|^2 \quad (70)$$

in full formal agreement with textbooks.

2.3 Quantum transitions in nuclear β -decay processes

The ongoing experimental and theoretical work in Nuclear and Particle Physics has by now initiated an exciting new era of fundamental physics. Random samples of the enormous output of publications in later years that are relevant to this paper are [9],[12],[13 -17],[18]).

An especially interesting and relevant feature of recent developments is the strong light they cast on the nature of quantum transitions in electroweak and QCD physics.

2.3.1 The time-dependent wavefunction

Good examples of contemporary interest is K-capture in highly ionized H-like atoms [18] and the physics of ultracold neutrons in traps [9]. We shall *reinterpret* this according to the philosophy sketched in this paper. Let us consider the decays

$${}_ZA_N + e^- \rightarrow {}_{Z-1}A_{N+1} + \nu_e \quad (71)$$

This is a K-capture process, meaning capture of an 1S atomic electron by a proton in a parent nucleus ${}_ZA_N$ with Z protons and N neutrons through

electroweak interactions that instantaneously replace a proton with a neutron in the daughter nucleus ${}_{Z-1}A_{N+1}$, together with an outgoing electron neutrino ν_e . This is accompanied by a sudden, but in this case rather small, change of the Coulomb field of the nucleus. An interesting feature of this kind of experiments is that the parent atom is supposed to be H-like, i.e. to be completely ionized except for a single bound electron, say in the 1S atomic state [18].

We shall consider this without any explicit or implicit reference to “measurements”, “probabilities”, “observations” or any other related standard concepts and terminology used in textbooks [2,3,4], where they are considered to be essential for self-consistency. As already discussed, this is not the case with the BP [2,3,4], where “measurements” of “observables” (implying actually no more than a permanent record of events stored somewhere and referring exclusively to a specific experimental set-up) are just special cases of quantum transition processes for beables [2,3,4]. Postulate 2 would link this to any specific available or predictable experimental results and, of course, to any textbook postdictions or predictions of such results.

Again, this is not the purpose of the present paper.

We imagine a movie showing the time development of a K-capture process occurring say at time $t=0$.

The Hamiltonian driving the evolution of the wave function is assumed to be

$$H = U_0 + H_0 + V_{eff} \quad (72)$$

U_0 represents the (infinite) Dirac vacuum energy. The next term is defined as

$$\begin{aligned} H_0 = & H_{nuc} - \frac{\hbar^2}{2m_e} \nabla_y^2(e) - Z\alpha \int d^3\vec{x} \frac{\varrho_{el}(\vec{x})}{|\vec{x} - \vec{y}|} + \\ & + \sum_{n=1}^3 [-i\hbar\vec{\alpha}(n) \cdot \vec{\nabla}_y(n) + \beta(n)m_n c^2] \end{aligned} \quad (73)$$

We shall reserve the notation $\{\vec{x}\}$ for space coordinates of the nucleons. The symbol $\vec{y}$ refers exclusively to the spatial coordinates of leptons. The first term on the rhs of (72) represents the nuclear Hamiltonian:

$$\begin{aligned} H_{nuc}\Phi_\alpha(\{\vec{x}, \vec{x}_{CM}\}; Z, N) = \\ = E_\alpha(Z, N)\Phi_\alpha(\{\vec{x}, \vec{x}_{CM}\}; Z, N) \end{aligned} \quad (74)$$

where

$$\vec{x}_{CM} \equiv \frac{M_P \sum_{i=1}^Z \vec{x}_{iP} + M_N \sum_{i=1}^N \vec{x}_{iN}}{ZM_P + NM_N} \quad (75)$$

are the CM coordinates of the nucleus defined in the atomic CM reference

system. The next two terms represent the Hamiltonian of the single atomic 1S electron with $\varrho_{el}(\vec{x})$ as the electric charge density profile of the nucleus. In the non-relativistic approximation

$$\begin{aligned} & \left[-\frac{\hbar^2}{2m_e} \nabla_y^2(e) - Z\alpha \int d^3\vec{x} \frac{\varrho_{el}(\vec{x})}{|\vec{x} - \vec{y}|} \right] \varphi_{1S\sigma}(\vec{y}) = \\ & = -|\varepsilon_{1S}| \varphi_{1S\sigma}(\vec{y}) \end{aligned} \quad (76)$$

The last term in definition (73) stands for the Dirac Hamiltonian of massive neutrinos, with the Bjorken-Drell conventions and definitions [20]. We need actually only the positive energy solutions $u_{n\vec{p}\sigma}(\vec{y})$:

$$\begin{aligned} & [-i\hbar\vec{\alpha}(n) \cdot \vec{\nabla}_y(n) + \beta(n)m_n c^2] u_{n\vec{p}\sigma}(\vec{y}) = \\ & = \sqrt{|\vec{p}|^2 c^2 + m_n^2 c^4} u_{n\vec{p}\sigma}(\vec{y}) \end{aligned} \quad (77)$$

The term V_{eff} in definition (72) is the electroweak interaction as originally suggested by the Standard Model [19] but adapted here (as an effective interaction) to the physical conditions inside nuclei [21]. As such it must be used with careful insight and according to standard ideas on modern renormalization techniques and effective field theories [21].

Assume that at time $t \leq 0$ the initial wavefunction of an individual H-like

ion at rest in its CM system is

$$\begin{aligned}\Psi_{I,1S\sigma}(\{\vec{x}, \vec{x}_M\}, \vec{y}; t) &\equiv \langle \{\vec{x}, \vec{x}_M\}, \vec{y} | I, 1S\sigma; t \rangle = \\ &= \Phi_I(\{\vec{x}, \vec{x}_M\}; Z, N) \varphi_{1S\sigma}(\vec{y}) e^{-iE_{I1S}t} \quad t \leq 0\end{aligned}\quad (78)$$

where $\Phi_I(\{\vec{x}, \vec{x}_M\}; Z, N)$ is the internal wavefunction of the parent nucleus. The total available energy is E_{I1S} :

$$E_{I1S} = U_0 + (M_I(Z, N) + m_e)c^2 - |\varepsilon(1S)| \quad (79)$$

where $M_I(Z, N)$ is the mass of the parent nucleus.

However, this state is not really a stationary state, because of the background weak couplings of quarks inside the nucleons to virtual flavour changing heavy electroweak bosons $W^\pm, Z^0$ [21]. This is represented by the (hermitean) last term V_{eff} on the rhs of the effective Hamiltonian (72) for allowed transitions [21]:

$$V_{eff}(\vec{x}, \vec{y}) = -\frac{G_F}{\sqrt{2}} \delta(\vec{x} - \vec{y}) (V_F(n) + V_{GT}(n)) \quad (80)$$

where $G_F = 8.7 \times 10^{-5} MeV.f m^2$ is the Fermi constant, $V_F(V_{GT})$ are related to the standard Fermi (Gamov-Teller) interaction operators.

Let us make the following ansatz for the time-dependent quantum state that should be good enough for sufficiently *weak interactions* [10]:

$$\begin{aligned}
& |I, 1S\sigma; t > = \\
& = \exp(-i(E_{I1S} - i\frac{\Gamma_{I1S}}{2})t) |I, 1S\sigma; t = 0 > \quad \Gamma_{I1S} > 0
\end{aligned} \tag{81}$$

It can be easily shown that [10]

$$\begin{aligned}
\Gamma_{I1S} &= 2\pi \sum_F \sum_{n, \sigma_n} \int d^3\vec{p}_n * \delta(E_{I1S} - E_{nF}(p_n)) * \\
& * | < F(-\vec{p}_n), n\vec{p}_n\sigma_n | V_{eff} | I, 1S\sigma > |^2
\end{aligned} \tag{82}$$

in agreement with Fermi's Golden Rule [10]. Furthermore

$$\begin{aligned}
E_{nF}(p_n) &= \\
&= U_0 + (M_F(Z - 1, N + 1)c^2 + \frac{|\vec{p}_n|^2}{2M_F} + \\
&+ \sqrt{|\vec{p}_n|^2 c^2 + m_n^2 c^4}
\end{aligned} \tag{83}$$

in obvious notation. The leptonic coordinates in the atomic CM system are related to $\vec{x}_{CM}$ by the definition

$$m_e \vec{y} + M_I \vec{x}_{CM} = 0 \quad (84)$$

in the initial state and

$$m_n \vec{y} + M_F \vec{x}_{CM} = 0 \quad (85)$$

in the final state.

We have used momentum conservation when writing down the expression for the recoil energy of the final nucleus.

The Bohm survival probability for an individual parent at any time $t > 0$ is defined to be

$$P_{I1S}(t) = \langle I, 1S\sigma; t | I, 1S\sigma; t \rangle = \exp(-\Gamma_{I1S} t) P_{I1S}(0) \quad (86)$$

This must satisfy the probability conservation law (25). So it does, to first order, since V_{eff} is by construction hermitean. Let us define as usual

$$\begin{aligned} \Psi_{I1S\sigma}(\{\vec{x}, \vec{x}_{CM}\}, \vec{y}; t) &= R_{I1S\sigma}(\{\vec{x}, \vec{x}_{CM}\}, \vec{y}; t) * \\ &= * \exp(i S_{I1S\sigma}(\{\vec{x}, \vec{x}_{CM}\}, \vec{y}; t)) \end{aligned} \quad (87)$$

where

$$\begin{aligned}
R_{I1S\sigma}(\{\vec{x}, \vec{x}_{CM}\}, \vec{y}; t) = \\
= \exp(-\frac{\Gamma_{I1S}}{2}t) |\Psi_{I,1S\sigma}(\{\vec{x}, \vec{x}_{CM}\}, \vec{y}, 0)|
\end{aligned} \tag{88}$$

$$P_{I1S}(t) = R_{I1S\sigma}^2(\{\vec{x}, \vec{x}_{CM}\}, \vec{y}; t) \tag{89}$$

Any built-in degeneracies of $\Psi_{I1S\sigma}(\{\vec{x}, \vec{x}_{CM}\}, \vec{y}; 0)$ would imply that its phase may not be zero.

2.3.2 Particle histories

In the context of the ontological BP under discussion a good question (though one that is never asked by any textbook!) could be: what are precisely the beable particle histories, i.e. the positions *and* velocities of those inside our decaying initial ion and the outgoing neutrino, at any time after $t=0$, until eventually some entirely new conditions - such as an observation! - occur [2]?

In the context of the BP of Quantum Theory, not only the question does make perfect sense, but can also be easily answered in quantitative detail, at least in perturbation theory as sketched above (even if, for the present, we are quite unable to plan, let alone carry out, any experiments based on such ideas!).

Especially interesting is any possibly “measurable quantity”, e.g. the mo-

momentum of the outgoing beable neutrino n or rather, the recoil energy of the daughter nucleus. The precise answer given by the BP is (section 1):

$$\begin{aligned} \vec{p}_n(\{\vec{X}(t), \vec{X}_{CM}(t)\}, \vec{Y}(t); t) = \\ = \{\vec{\nabla}_{\vec{y}} S_{I1S\sigma}(\{\vec{x}, \vec{x}_{CM}(t)\}, \vec{y}; t)\}_{\{\vec{x}\}\vec{x}_{CM}, \vec{y}=\{\vec{X}(t)\}, \vec{X}_{CM}(t), \vec{Y}(t)} \end{aligned} \quad (90)$$

Immediate confrontation with existing experimental material on this question is however possible (Postulate 2 ,section 1) if one first carries out an appropriate stochastic averaging over all initial positions and velocities of the (entangled) beable nucleons and leptons initially involved :

$$\begin{aligned} < \vec{p}_n(\{\vec{X}(t), \vec{X}_{CM}(t)\}, \vec{Y}(t); t) >_{EA} = \int \int \int [R_{I1S\sigma}^2(\{\vec{x}\}\vec{x}_{CM}, \vec{y}; t)] * \\ * \vec{\nabla}_y(n) S_{I1S\sigma}(\{\vec{x}\}\vec{x}_{CM}, \vec{y}; t) \}_{\{\vec{x}\}\vec{x}_{CM}, \vec{y}=\{\vec{X}(t)\}\vec{X}_{CM}(t), \vec{Y}(t)} \end{aligned} \quad (91)$$

The integrations are to be carried out over all histories of all participating beable particles. As mentioned in section 2, this is guaranteed to be in numerical agreement with the textbook answer at $t=0$. As also mentioned in section 2, it will then be true at any time :

$$< (-i\vec{\nabla}_y(n)) > =$$

$$\begin{aligned}
&= \frac{1}{2} \int \int [\Psi_{I1S\sigma}^* (-i\vec{\nabla}_y(n) \Psi_{I1S\sigma}) + \Psi_{I1S\sigma} (-i\vec{\nabla}_y(n) \Psi_{I1S\sigma})^*] \equiv \\
&\equiv \langle \vec{p}_n(\{\vec{X}(t)\} \vec{X}_{CM}(t), \vec{Y}(t); t) \rangle_{EA} \quad (92)
\end{aligned}$$

We emphasize that the word “time” means here literally “time taken by a particle, say the neutrino n , to be guided from its position $\vec{X}(t)$ (and velocity $\vec{V}(t)$) to position $\vec{X}'(t')$ (and velocity $\vec{V}'(t')$)”. If a “distance” L is “measured” from the neutrino position to the CM of the parent atom, then at time $t > 0$ the distance L can be traded with time t elapsed because to sufficient accuracy $t = L/c$. Hence, the time label t in (91) can be replaced by the space label L , and so all talk about a possibly vague “time” can be avoided [22]. The point being, “time” here is directly linked to the physical motions of material particles.

3 Summary and conclusions

The ontological formulation of non-relativistic Quantum Mechanics, originally initiated by Louis de Broglie in the twenties but independently recreated and reformulated in the fifties and sixties by D. Bohm and his co-workers, is briefly presented and its explanatory powers are further illustrated with examples from new fields of research in contemporary Nuclear and Particle

Physics, fields that were not mature during the lifetime of those pioneers.

It is argued that the reformulation in question has indeed the capability of not only by-passing all the so-called paradoxes and ambiguities characterizing the official (textbook) epistemological formulations of Quantum Theory, but more importantly, lead to genuine testable *physical explanations* (and not only to highly abstract mathematical representations) of actual physical phenomena, at the quantum level of accuracy. This is achieved without any need of arbitrary breaks of continuity between the so-called “classical” and “quantal” modes of description that is so vital for all textbook formulations. Furthermore, this novel ontological formulation can lead naturally to proposals for new type of experimentation, thus in principle greatly extending the predictive powers of present day Quantum Theory. One can thus open up for new ways of thinking and trying fresh approaches to the age-old fundamental problems of Relativistic Quantum Field Theory and Theoretical Cosmology.

References

- [1] P.A.M.Dirac “Lectures on Quantum Field Theory” (Academic Press inc.,New York,USA,1966)
- [2] D.Bohm and B.J.Hiley ”The undivided universe” (Routledge, London, England, 1993)

- [3] P. Holland "The quantum theory of motion" (Cambridge University Press, Cambridge, 1977)
- [4] J.S.Bell "Speakable and unspeakable in quantum mechanics" (Cambridge University Press, Cambridge, UK,1996)
- [5] J. Passon, quant-ph/0611032
- [6] H.O.U. Fynbo et al, Phys.Rev.Let.,91(2003)082502
- [7] S.Ali and A.R.Bodmer Nucl.Phys.80(1966)99-112
- [8] H.O.U. Fynbo et al, Nucl. PhysA318(2003)541c-543c
- [9] J.S.Nico and W.M.Snow, Ann.Rev.Nucl.Part.Sci.55(2005) 27-69
- [10] J.J.Sakurai"Modern Quantum Mechanics"(Addison - Wesley Publ Co,Inc,New York, USA, 1985)
- [11] Ray d'Inverno "Introducing Einstein's relativity" (Oxford University Press, UK, 1996)
- [12] B.Kayser, hep-ph/0804.1497
- [13] T.D.Lee, hep-ph/0605017
- [14] R.Friedberg and T.D.Lee, hep-ph/060607
- [15] R.Friedberg and T.D.Lee, hep-ph/709.1526
- [16] R.Friedberg and T.D.Lee, hep-ph/0705.4156

- [17] R.Friedberg and T.D.Lee, hep-ph/...09.3633
- [18] Litvinov et al, nucl-ex/08012079
- [19] S. Weinberg “The Quantum Theory of Fields”, vol. II (Cambridge University Press, Cambridge, UK, 1996)
- [20] J. Bjorken and S. Drell “Relativistic Quantum Fields” (McGraw Hill Book Company, New York, USA, 1965)
- [21] P.J. Siemens and A. Jensen ”Elements of nuclei” (Addison Wesley Publishing Co, Inc, New York, USA, 1987)
- [22] H. Lipkin, hep-ph/9901399